

Drexel

# Beau-Ron D

Soluble Borate Micronutrient

(Disodium octaborate tetrahydrate)

### GUARANTEED ANALYSIS:

Boron (B)<sup>1</sup> ..... 20.96%  
<sup>1</sup>Derived from Sodium borate.  
 4.8 pounds of BEAU-RON D provides 1 pound elemental Boron.  
 Ingredients exempt from tolerance under 40 CFR 180.1001

**KEEP OUT OF REACH OF CHILDREN  
 CAUTION**

Read Entire Label Before Using This Product

Net Weight: 10 lb (4.54 kg)

### PRECAUTIONARY STATEMENTS

**Hazards to Humans and Domestic Animals**  
 Harmful if swallowed or inhaled. Avoid contact with eyes, skin and clothing. Causes eye and skin irritations. In case of contact, wash the affected areas with water immediately. Call a doctor if irritation persists.

### GENERAL INFORMATION

BEAU-RON D soluble borate is for fluid fertilizer nutrient spray and dusts. The total amount of this product applied over the growing season should not exceed the total recommended for your area (typically between 1 to 2 pounds elemental boron per acre). Boron deficiency symptoms can usually be corrected in most crops in 10 to 14 days. Do not exceed one-half pound of elemental boron per acre per application.

### DIRECTIONS FOR USE

This product is recommended for aerial, concentrate and dilute application to field, vegetable, tree and nut crops. Rates and timing of applications are dependent on local conditions and should be made only in conjunction with the recommendation of the Local authority. Soil and/or plant tissue analysis should be obtained and the results used as a guide. Thorough coverage of this product is necessary for best results and sufficient water should be used to assure uniform application.

**SOIL APPLICATION:** This product can be sprayed directly on the soil with a boom sprayer if a soil application of boron is desired. Application rates are the same as for a foliar spray.

**FOLIAR SPRAYS AND DUSTS:** This product applied to the foliage of growing plants at rates of 4.8 to 9.6 pounds per acre, will correct boron deficiency in most crops. It is compatible with most insecticides and fungicides and can be included in many established applications being made for the control of insects and fungus diseases. Spray applications may be made with either concentrate or high volume sprayers.

**LIQUID FERTILIZERS:** This product is readily soluble in water and makes an efficient borate additive for most liquid fertilizers. It is superior to borax as it is more readily soluble and causes minimum changes in crystallization temperatures.

**NOTE:** Rates and timing of boron applications are dependent on local conditions and should be made as a result of soil and/or plant tissue analysis and in conjunction with your Local Agricultural Authority.

### CROP AND RATE RECOMMENDATIONS

Field Crops	Pounds per Acre
Alfalfa, Clover	2.4 to 4.8
Coffee	Consult Local Usage
Corn, Cotton	1.2 to 2.4
Lentils, Peas (field)	1.2 to 2.4
Peanuts	1.2 to 2.4
Rape (Canola) Sorghum	1.2 to 2.4
Soybeans	1.2 to 2.4
Sugarbeet <sup>1</sup> , Sunflower <sup>1</sup>	2.4 to 4.8
Wheat Barley Oats	1.2 to 2.4

Fruit and Nut Crops	Pounds per Acre
Almonds	2.4 to 4.8
Apple, Peach Pear <sup>2</sup>	2.4 to 4.8
Apricot <sup>2</sup>	7.7 to 9.6
Avocado	Soil and/or Plant Analysis
Cherries	Soil and/or Plant Analysis
Citrus <sup>2</sup>	1.2 to 3.8
Fig	Soil and/or Plant Analysis
Grapes <sup>2</sup>	3.8 to 6.8
Pecan	Soil and/or Plant Analysis
Plums and Prunes <sup>2</sup>	2.4 to 9.6
Strawberries	2.4 to 4.8
Walnuts <sup>2</sup>	7.2 to 9.6

Vegetable Crops	Pounds per Acre
Artichoke, Beans	Soil and/or Plant Analysis
Beets (red table)	2.4 to 4.8
Broccoli	3.8 to 9.6
Brussels sprouts Cabbage	1.2 to 7.7
Carrots	1.2 to 2.4
Cauliflower	2.4 to 9.6
Celery, Lettuce Onions	1.2 to 3.8
Potatoes	1.2 to 4.8
Pumpkin, Sweet potatoes	Soil and/or Plant Analysis
Radishes, Spinach	1.2 to 3.8
Tomatoes, Turnips	1.2 to 3.8

<sup>1</sup>Apply up to 2.4 pounds of this product when Sugarbeets and Sunflowers are at least six (6) inches high. Apply up to 2.4 pounds approximately thirty (30) days later.

<sup>2</sup>Care should be taken when applying this product to these crops. Excessive amounts of boron may cause crop damage.

The following chart can be used as a guide for applying the correct rates of this product:

CONVERSION CHART	
Pounds of Boron (B) to be Applied per Acre	Amount of This Product Needed per Acre
0.10	0.50 lbs.
0.20	1.00 lbs.
0.50	2.40 lbs.
0.75	3.60 lbs.
1.00	4.80 lbs.
1.50	7.20 lbs.
2.00	9.60 lbs.

Manufactured By:

**Drexel Chemical Company**

P.O. BOX 13327, MEMPHIS, TN 38113-0327

SINCE 1972

### **STORAGE AND DISPOSAL**

Do not contaminate water, food or feed by storage and disposal.  
**PRODUCT DISPOSAL:** Follow all governmental requirements for the disposal of spray solutions.

**CONTAINER DISPOSAL:** Dispose of empty container according to approved Local practices. Contact County Agricultural Commissioner for disposal information. Do not reuse container.

### **WARRANTY—CONDITIONS OF SALE**

OUR RECOMMENDATIONS FOR USE of this product are based upon tests believed reliable. Follow directions carefully. Timing and method of application, weather and crop conditions, mixtures with other chemicals not specifically recommended and other influencing factors in the use of this product are beyond the control of the Seller. Buyer assumes all risks of use, storage and handling of this material not in strict accordance with directions given herewith.

In no case shall the Manufacturer or the Seller be liable for consequential, special or indirect damages resulting from the use or handling of this product when such use and/or handling is not in strict accordance with directions given herewith. The foregoing is a condition of sale by the Seller and is accepted as such by the Buyer.