

Caution: This product contains BORON and MOLYBDENUM. Determine plant sensitivity prior to use. Plants containing excess MOLYBDENUM are toxic to livestock. Crops sensitivity to either or these two elements may be injured



ENERGY BOOSTER™



0-0-7

GUARANTEED ANALYSIS

Soluble Potash (K ₂ O).....	7%
Magnesium (Mg) Total.....	3.25%
3.25% Water Soluble Magnesium (Mg)	
1.0% Chelated Magnesium (Mg)	
Sulfur (S) Total.....	2.45%
2.45% Combined Sulfur (S)	
Boron (B).....	0.20%
Cobalt (Co).....	0.03%
0.03% Chelated Cobalt	
Copper (Cu).....	0.90%
0.90% Chelated Copper (Cu))	
Iron (Fe)	2.40%
2.40% Chelated Iron (Fe)	
Manganese (Mn)	2.40%
2.40% Chelated Manganese (Mn)	
Molybdenum (Mo).....	0.06%
Zinc (Zn)	0.90%
0.90% Chelated Zinc (Zn)	

Derived From: Potassium Sulfate, Kelp, Carboxylic Acid, Magnesium EDTA, Iron EDTA, Manganese EDTA, Zinc EDTA, Cobalt EDTA, Magnesium Sulfate, Sodium Borate and Ammonium Molybdate.

Chelating agent is EDTA (ethylenediaminetetraacetic acid).

Chloride (Cl) maximum..... 0.12%

Potential Acidity is equivalent to 469 lbs. Calcium Carbonate (CaCO₃) per ton.

MANUFACTURED FOR
CNI AgriMinerals LLC.
 P.O. BOX 3706
 Albany, GA. 31706 USA

GENERAL INFORMATION

ENERGY BOOSTER is formulated to be compatible with Glyphosate and other selective and non-selective herbicides. Research conducted with tank mixes of the chelating agents used in **ENERGY BOOSTER** helped to maintain the efficacy of the herbicides. This product gives best results when crops have adequate supplies of water and major nutrients, and are not under stress for other reasons. Conditions which are responsible for one particular deficiency can also induce deficiencies of other trace elements.

ENERGY BOOSTER is particularly suitable for foliar application and for general soil applications. Foliar fertilization is intended as a supplement to a regular fertilization program and will not, by itself, provide all the nutrients normally required by agricultural crops. Plants need only very small amounts of micronutrients for maximum growth.

CROPS SUSCEPTIBLE TO NUTRITIONAL MICRONUTRIENT DEFICIENCY

ENERGY BOOSTER is recommended for crops such as alfalfa, almonds, avocados, apples, broccoli, cabbage, canola, cauliflower, carrots, celery, citrus, corn, cotton, grapes, lettuce, milo, melons, nectarines, rice, pears, peaches, pecans, peppers, plums, prunes, potatoes, peanuts, sorghum, soybeans, sugar beets, sweet corn, sugar cane, strawberries, tomatoes, turnips, walnuts, watermelons, wheat and most other crops.

MIXING

ENERGY BOOSTER is soluble in water. When mixing; do NOT pre-mix a small amount of water before adding to the spray tank. **ENERGY BOOSTER** should be added slowly to the majority of the spray water volume while it is being agitated. Agitation should continue for a minute or two after the addition of **ENERGY BOOSTER**. If pesticides are to be used in the spray mix, add them after the **ENERGY BOOSTER** has dissolved.

RECOMMENDED RATES

Soil: Apply 1 lb. per acre of **ENERGY BOOSTER** in combination with starter or pop-up fertilizers. **ENERGY BOOSTER** can also be applied at 1 lb per acre broadcast with pre-plant or pre-emergence herbicide application.

Foliar: Apply 1 lb per acre of **ENERGY BOOSTER** in a minimum of 20 gallons of water per acre. Applications made in lower volumes of water, especially if the crop is under stress, may cause injury. Therefore, use no more than ¼ lb of **ENERGY BOOSTER** per 5 gallons of water used per acre.

Aerial application: Apply at the same use rate of ¼ lb per 5 gallons of water used per acre. **DO NOT** exceed 1 lb. per acre per application even if exceeding 20 gallons of water per acre.

CAUTION: Do not apply more than 1 lb. per acre of **ENERGY BOOSTER** per foliar application as crop injury may occur. Multiple applications should be made at no less than 10 to 14 day intervals to reduce the potential for crop injury. Wash hands with soap and water before meals and after work. Store in the original container tightly closed, in a dry place. Store away from children, pets, livestock and foodstuffs.

Use this product in accordance with good agronomic practices, which include utilizing proven spray equipment set for proper coverage. Do not make applications when temperatures are too hot, as crop damage may occur. Applications should be made at temperature levels and when other environmental conditions in your area are such that your experience indicates the application will be compatible and will accomplish the desired result.

Notice of Warranty: **CNI AgriMinerals LLC** warrants that this product conforms to the chemical description on the label thereof and is reasonably fit for purposes stated on such label only when used in accordance with directions under normal use conditions. It is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness, or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of **CNI AgriMinerals LLC**. In no case shall **CNI AgriMinerals LLC** be liable for consequential, special or indirect damages resulting from the use or handling of this product. All such risks shall be assumed by the buyer. **CNI AgriMinerals LLC** MAKES NO WARRANTIES OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS STATED ABOVE.

Information concerning the raw materials composing this product can be obtained by writing to **CNI AgriMinerals LLC**, P.O. Box 3706, Albany, GA. 31706. Please refer to the batch number found on this container.

KEEP OUT OF REACH OF CHILDREN
CAUTION

Net Weight 25 Lb (11.34 kg)